

2014-2015 Season (9/28/2014 – 10/3/2015)

**Synopsis:**


Arizona identified the first locally-acquired influenza case of the 2014-2015 season during week 45, a child infected with influenza A (H3) as confirmed by PCR at the Arizona State Public Health Laboratory. Influenza activity otherwise continued to be low. Arizona reported Sporadic activity for week 45.

**Influenza activity highlights:**

- The first locally-acquired influenza case of the 2014-2015 season was identified during week 45.
- 18 laboratory-confirmed cases of influenza were reported in the past week, from 4 counties. 21 cases have been reported this season, with laboratory-confirmed cases identified in 4 counties. The first three cases reported were not confirmed to be locally-acquired.
- 14 (67%) reports this season are influenza A, 6 (29%) are influenza B, and 1 (5%) is of unknown type.
- 13 (42%) of 31 specimens tested positive for influenza at ASPHL last week; 12 influenza A (H3) viruses and 1 influenza B virus.
- Influenza-like illness activity at sentinel providers was below Arizona’s threshold in week 44.
- Influenza-like illness activity at sentinel schools decreased in week 45.
- The cases included in this report represent a small proportion of the true number of cases of influenza. Many people do not visit the doctor when ill and doctors should not be expected to run tests on all patients exhibiting influenza-like symptoms.

**Laboratory-Confirmed Influenza Activity by Season [2009-2015]**

Positive influenza tests are reported to ADHS. Many types of tests are included in the numbers below: rapid antigen tests, direct fluorescent antigen tests, viral culture, and molecular testing.



## Reported Laboratory-Confirmed Cases Compared to Last Week and Last Season

	Cumulative Season Total	Current Week Total
<b>2014-2015</b>	21	18
<b>2013-2014</b>	91	29
<b>5 year average</b>	30	11
<b>% increase, compared to 2013-2014 season</b>	-77%	-38%
<b>% increase, compared to a typical flu season</b>	-31%	61%
<b>% increase, compared to last week</b>	600%	1,700%


\* Mean of past 5 years, excludes 2009-2010 pandemic season

### Arizona Influenza Activity Levels *(see definitions at the end of this report)*

	2014-2015	2013-2014	2012-2013	2011-2012	2010-2011
<b>This Week</b>	Sporadic	Sporadic	Sporadic	No Activity	Sporadic
<b>Last Week</b>	Sporadic	Sporadic	Sporadic	No Activity	Sporadic
<b>Date First Case Confirmed, no travel</b>	Nov. 3, 2014	Oct. 4, 2013	Oct. 30, 2012	Dec. 14, 2011	Sept. 28, 2010
<b>Weeks with Widespread Activity</b>	No widespread activity yet	Weeks 3-8	Weeks 1-7	Weeks 11 – 14	Weeks 52 – 10

### Laboratory-Confirmed Cases Reported, by County, 2014-2015 Influenza Season (Includes ALL reported lab-confirmed flu reports, regardless of subtype)

County	2014-2015 Season	Past Three Weeks	Last Week
Apache	0	0	0
Cochise	0	0	0
Coconino	0	0	0
Gila	0	0	0
Graham	0	0	0
Greenlee	0	0	0
La Paz	0	0	0
Maricopa	7	6	6
Mohave	0	0	0
Navajo	0	0	0
Pima	11	10	9
Pinal	2	2	2
Santa Cruz	0	0	0
Yavapai	1	1	1
Yuma	0	0	0
<b>Total</b>	21	19	18



**Key:**

- \* = Any activity reported this season
- = Activity reported in the previous three weeks
- = No activity reported in the previous three weeks

**Age of Reported Influenza Cases**

The age groups most affected by influenza vary somewhat season-to-season, depending in part on the circulating influenza types and subtypes and any existing immunity in the community. Variations in age groups of reported influenza cases can also be caused by differences in laboratory testing and reporting practices year-to-year.

**Age Group of Reported Influenza Cases, 2011-2012 through 2014-2015 Seasons**

Age Group	2014-2015 Season (N=21)	2013-2014 Season (N=12,484)	2012-2013 Season (N=11,301)	2011-2012 Season (N=4,004)
0 to 4 years	2 (10%)	2,329 (19%)	2,114 (19%)	750 (19%)
5 to 18 years	7 (33%)	2,802 (22%)	3,013 (27%)	1,053 (26%)
19 to 49 years	4 (19%)	4,487 (36%)	3,107 (27%)	1,352 (34%)
50 to 64 years	4 (19%)	1,566 (13%)	1,156 (10%)	400 (10%)
65 years or older	4 (19%)	1,205 (10%)	1,799 (16%)	436 (11%)
Unknown age	0 (0%)	95 (1%)	112 (1%)	13 (0.3%)


**Age Group of Reported Influenza Cases by Type, 2014-2015 Season**

Age Group	All Confirmed Cases (N=21)	Influenza A (N=14)	Influenza B (N=6)	Unknown Type (N=1)
0 to 4 years	2 (10%)	0 (0%)	2 (33%)	0 (0%)
5 to 18 years	7 (33%)	7 (50%)	0 (0%)	0 (0%)
19 to 49 years	4 (19%)	2 (14%)	2 (33%)	0 (0%)
50 to 64 years	4 (19%)	4 (29%)	0 (0%)	0 (0%)
65 years or older	4 (19%)	1 (7%)	2 (33%)	1 (100%)
Unknown age	0 (0%)	0 (0%)	0 (0%)	0 (0%)

## Influenza Types and Subtypes

There are two main types of influenza – Type A and Type B – that cause illness in people. Influenza A viruses can be further divided into subtypes such as A (H1), or A (H3). While most tests can distinguish between influenza A and B, only specialized testing such as that done at the State Public Health Laboratory and a few other labs around the state can differentiate subtypes. Viral culture or molecular testing (reverse transcriptase polymerase chain reaction or RT-PCR) are the methods used to identify subtypes; knowing the type and subtype of the influenza viruses circulating can help health professionals make the best treatment and vaccination decisions.

### Influenza Type, from all tests reported




### Influenza Type, by Season

	2014-2015 Season Number	2014-2015 Season Percent	2013-2014 Number (Percent)	2012-2013 Number (Percent)	2011-2012 Number (Percent)
<b>Total</b>	21	100%	11,780 (100%)	11,306 (100%)	4,004 (100%)
<b>Influenza A</b>	14	67%	9,355 (79%)	8,059 (71%)	2,820 (70%)
<b>Influenza B</b>	6	28%	2,229 (19%)	2,951 (26%)	1,078 (27%)
<b>Unknown</b>	1	5%	196 (2%)	296 (3%)	106 (3%)

## Influenza Subtype

### Data from the Arizona State Public Health Laboratory (ASPHL)

- In the past week, 13 (42%) of 31 specimens tested positive for influenza at ASPHL.
- 12 are influenza A (H3) viruses and 1 is an influenza B virus



### Influenza subtype, by season, from any laboratories performing culture or RT-PCR


	2014-2015 Season Number	2014-2015 Season Percent	2013-2014 Number (Percent)	2012-2013 Number (Percent)	2011-2012 Number (Percent)
<b>Influenza Subtypes</b>	8	100%	3,307 (100%)	3,243 (100%)	1,564 (100%)
Influenza A (H1N1)pdm09	0	0%	1,467 (44%)	79 (2%)	594 (38%)
Influenza A (H3)	2	25%	132 (4%)	1,588 (49%)	351 (22%)
Influenza A (Unsubtyped)	2	25%	1,115 (34%)	829 (26%)	255 (16%)
Influenza B	4	50%	593 (18%)	747 (23%)	364 (23%)

## Severe Acute Respiratory Infections (SARI) Surveillance

Severe acute respiratory infections (SARI) surveillance is currently conducted at selected hospitals in counties along the Mexican border by the Office of Border Health's Border Infectious Disease Surveillance (BIDS) program. SARI is defined as a hospital admission with a fever of at least 100°F plus either a cough or a sore throat. This surveillance facilitates the detection of circulating influenza viruses and allows us to monitor various causes of morbidity and mortality among inpatients with SARI.

SARI cases are tested using an RT-PCR viral panel that detects: influenza A (H1) seasonal subtype, A (H3) seasonal subtype, A (H1N1) pdm09, and B; respiratory syncytial virus A and B; parainfluenza virus 1, 2, 3, and 4; human metapneumoviruses A/B; rhinovirus; adenovirus (B, C, and E); and coronavirus (NL63, HKU1, 229E, and OC43). Bacterial testing is conducted for *Mycoplasma pneumoniae*, *Chlamydia pneumoniae*, *Legionella pneumophila* and *Bordetella pertussis* if a bacterial specimen is available for off-site testing. However, if a bacterial specimen is not submitted, patient chart reviews are conducted to find additional bacterial testing results performed on site.


10 suspect SARI cases have been identified this season. Parainfluenza has been detected in one case.



## Influenza-Like Illness (ILI) Surveillance from Sentinel Outpatient Providers

ILI is defined as a fever of at least 100°F plus either a cough or a sore throat. In weeks when a relatively low number of enrolled facilities report data, the ILI proportion may not be as representative of Arizona activity as for other weeks. The state ILI baseline is 1.1% and the epidemic threshold is 1.9%\*.

	Week 44	Week 43
Proportion of patient visits to sentinel providers for ILI	1.0%	1.0%
Comparison to epidemic threshold*	Below threshold	Below threshold
Intensity level <i>(see definitions at the end of report)</i>	Minimal	Minimal




\*Note: The baseline is defined as the mean of the state ILI% in weeks in the 2011-2014 flu seasons in which two or more consecutive weeks each accounted for less than 2% of the season's total number of specimens testing positive for influenza at the Arizona State Public Health Laboratory. The epidemic threshold is defined as the mean plus two standard deviations.

## School Surveillance for Influenza-Like Illness (ILI)

School nurses in approximately 140 Arizona schools around the state use a specific computer program (the Child Health Indicator Program) for electronic management of student health records. The graph presents the weekly trend of ILI syndromes reported among students during the past four school years. School nurse encounters are not diagnosed cases of communicable diseases but are based on the nursing codes that school nurses enter to track student conditions. Also, the numbers in the graph are only from schools that used CHIP during the school year.

### Influenza-Like Illness Per School Enrollment in Arizona (per 100,000)



## **Glossary of Key Terms:**

2014-2015 Influenza Season – The season is defined by surveillance weeks. The first day of the 2014-2015 influenza season was September 28<sup>th</sup>, 2014, or week 40 and the 2014-2015 surveillance season will continue through October 3<sup>rd</sup>, 2015, or week 39.

Regions – Regions in Arizona are defined by county: Central (Gila, Maricopa, Pinal); Northern (Apache, Coconino, Navajo, Yavapai); Southern (Cochise, Graham, Greenlee, Pima, Santa Cruz); Western (La Paz, Mohave, Yuma)

Activity Levels: Indicator of the geographic spread of influenza activity, reported to CDC by all states each week.

Widespread: Increased influenza-like illness from sentinel providers (ILI) in three or more regions and large numbers of laboratory-confirmed influenza cases in those regions.

Regional: Increased ILI in two regions and elevated numbers of laboratory-confirmed influenza cases in those regions.

Local: Increased ILI in one region and elevated numbers of laboratory-confirmed influenza cases in that region.

Sporadic: No increase in ILI activity and only isolated laboratory-confirmed influenza cases.

No Activity: No increase in ILI activity and no laboratory-confirmed influenza cases.

Intensity Levels: Intensity levels are based on the percent of outpatient visits in a state due to ILI and are compared to the average percent of ILI visits that occur during spring and fall weeks with little or no influenza virus circulation. Intensity levels range from minimal, corresponding to ILI activity from outpatient clinics being below the average, to intense, which would correspond to ILI activity from outpatient clinics being much higher than average.