

2015-2016 Season (10/4/2015 – 10/1/2016)

Synopsis:


Arizona has identified the first locally-acquired influenza case of the 2015-2016 season. During the past week, influenza activity continued to be low. Arizona reported Sporadic activity for week 40.

Influenza activity highlights:

- The first locally-acquired influenza cases of the 2015-2016 season have been identified, with confirmation at the Arizona State Public Health Laboratory.
- 7 laboratory-confirmed cases of influenza were reported in the past week, from two counties.
- No specimens tested positive for influenza at the Arizona State Public Health Laboratory in the past week.
- Influenza-like illness activity at sentinel providers was below Arizona’s threshold in week 39.
- The cases included in this report represent a small proportion of the true number of cases of influenza. Many people do not visit the doctor when ill and doctors should not be expected to run tests on all patients exhibiting influenza-like symptoms.

Laboratory-Confirmed Influenza Activity by Season [2009-2016]

Positive influenza tests are reported to ADHS. Many types of tests are included in the numbers below: rapid antigen tests, direct fluorescent antigen tests, viral culture, and molecular testing.



Reported Laboratory-Confirmed Cases Compared to Last Week and Last Season

| | Cumulative Season Total | Current Week Total |
|---|-------------------------|--------------------|
| 2015-2016 | 7 | 7 |
| 2014-2015 | 1 | 1 |
| 5 year average | 1 | 1 |
| % increase, compared to 2014-2015 season | 600% | 600% |
| % increase, compared to a typical flu season | 483% | 483% |
| % increase, compared to last week | N/A | N/A |


Arizona Influenza Activity Levels *(see definitions at the end of this report)*

| | 2015-2016 | 2014-2015 | 2013-2014 | 2012-2013 | 2011-2012 |
|---|----------------------------|--------------|--------------|---------------|---------------|
| This Week | Sporadic | No Activity | Sporadic | Sporadic | Sporadic |
| Last Week | Sporadic | Sporadic | No Activity | No Activity | No Activity |
| Date First Case Confirmed, no travel | Oct. 4, 2015 | Nov. 3, 2014 | Oct. 4, 2013 | Oct. 30, 2012 | Dec. 14, 2011 |
| Weeks with Widespread Activity | No widespread activity yet | Weeks 1-6 | Weeks 3-8 | Weeks 1-7 | Weeks 11 – 14 |

Laboratory-Confirmed Cases Reported, by County, 2015-2016 Influenza Season

(Includes ALL reported lab-confirmed flu reports, regardless of subtype)

| County | 2015-2016 Season | Past Three Weeks | Last Week |
|--------------|------------------|------------------|-----------|
| Apache | 0 | 0 | 0 |
| Cochise | 0 | 0 | 0 |
| Coconino | 0 | 0 | 0 |
| Gila | 0 | 0 | 0 |
| Graham | 0 | 0 | 0 |
| Greenlee | 0 | 0 | 0 |
| La Paz | 0 | 0 | 0 |
| Maricopa | 5 | 5 | 5 |
| Mohave | 0 | 0 | 0 |
| Navajo | 0 | 0 | 0 |
| Pima | 0 | 0 | 0 |
| Pinal | 0 | 0 | 0 |
| Santa Cruz | 0 | 0 | 0 |
| Yavapai | 2 | 2 | 2 |
| Yuma | 0 | 0 | 0 |
| Total | 7 | 7 | 7 |



Key:

- * = Any activity reported this season
- = Activity reported in the previous three weeks
- = No activity reported in the previous three weeks

Age of Reported Influenza Cases

The age groups most affected by influenza vary somewhat season-to-season, depending in part on the circulating influenza types and subtypes and any existing immunity in the community. Variations in age groups of reported influenza cases can also be caused by differences in laboratory testing and reporting practices year-to-year.

Age Group of Reported Influenza Cases, 2012-2013 through 2015-2016 Seasons

| Age Group | 2015-2016 Season (N=7) | 2014-2015 Season (N=12,580) | 2013-2014 Season (N=12,484) | 2012-2013 Season (N=11,301) |
|-------------------|---------------------------|--------------------------------|--------------------------------|--------------------------------|
| 0 to 4 years | 0 (0%) | 2,153 (17%) | 2,329 (19%) | 2,114 (19%) |
| 5 to 18 years | 0 (0%) | 3,365 (27%) | 2,802 (22%) | 3,013 (27%) |
| 19 to 49 years | 0 (0%) | 3,039 (24%) | 4,487 (36%) | 3,107 (27%) |
| 50 to 64 years | 1 (14%) | 1,223 (10%) | 1,566 (13%) | 1,156 (10%) |
| 65 years or older | 6 (86%) | 2,659 (21%) | 1,205 (10%) | 1,799 (16%) |
| Unknown age | 0 (0%) | 141 (1%) | 95 (1%) | 112 (1%) |


Age Group of Reported Influenza Cases by Type, 2015-2016 Season

| Age Group | All Confirmed Cases (N=7) | Influenza A (N=6) | Influenza B (N=1) | Unknown Type (N=0) |
|-------------------|------------------------------|----------------------|----------------------|-----------------------|
| 0 to 4 years | 0 (0%) | 0 (0%) | 0 (0%) | 0 (0%) |
| 5 to 18 years | 0 (0%) | 0 (0%) | 0 (0%) | 0 (0%) |
| 19 to 49 years | 0 (0%) | 0 (0%) | 0 (0%) | 0 (0%) |
| 50 to 64 years | 1 (14%) | 1 (17%) | 0 (0%) | 0 (0%) |
| 65 years or older | 6 (86%) | 5 (83%) | 1 (100%) | 0 (0%) |
| Unknown age | 0 (0%) | 0 (0%) | 0 (0%) | 0 (0%) |

Influenza Types and Subtypes

There are two main types of influenza – Type A and Type B – that cause illness in people. Influenza A viruses can be further divided into subtypes such as A (H1), or A (H3). While most tests can distinguish between influenza A and B, only specialized testing such as that done at the State Public Health Laboratory and a few other labs around the state can differentiate subtypes. Viral culture or molecular testing (reverse transcriptase polymerase chain reaction or RT-PCR) are the methods used to identify subtypes; knowing the type and subtype of the influenza viruses circulating can help health professionals make the best treatment and vaccination decisions.

Influenza Type, from all tests reported



Influenza Type, by Season

| | 2015-2016 Season Number | 2015-2016 Season Percent | 2014-2015 Number (Percent) | 2013-2014 Number (Percent) | 2012-2013 Number (Percent) |
|--------------------|-------------------------------|--------------------------------|----------------------------------|----------------------------------|----------------------------------|
| Total | 7 | 100% | 12,592 (100%) | 11,780 (100%) | 11,306 (100%) |
| Influenza A | 6 | 86% | 11,015 (87%) | 9,355 (79%) | 8,059 (71%) |
| Influenza B | 1 | 14% | 1,424 (11) | 2,229 (19%) | 2,951 (26%) |
| Unknown | 0 | 0% | 153 (1%) | 196 (2%) | 296 (3%) |

Influenza Subtype

Data from the Arizona State Public Health Laboratory (ASPHL)

- No specimens tested positive for influenza at the Arizona State Public Health Laboratory in the past week.


Influenza subtype, by season, from any laboratories performing culture or RT-PCR

| | 2015-2016 Season Number | 2015-2016 Season Percent | 2014-2015 Number (Percent) | 2013-2014 Number (Percent) | 2012-2013 Number (Percent) |
|---------------------------|-------------------------------|--------------------------------|----------------------------------|----------------------------------|----------------------------------|
| Influenza Subtypes | 3 | 100% | 2,955 (100%) | 3,307 (100%) | 3,243 (100%) |
| Influenza A (H1N1)pdm09 | 0 | 0% | 5 (0.1%) | 1,467 (44%) | 79 (2%) |
| Influenza A (H3) | 1 | 33% | 2,114 (72%) | 132 (4%) | 1,588 (49%) |
| Influenza A (Unsubtyped) | 2 | 67% | 488 (17%) | 1,115 (34%) | 829 (26%) |
| Influenza B | 0 | 0% | 348 (12%) | 593 (18%) | 747 (23%) |

Influenza-Like Illness (ILI) Surveillance from Sentinel Outpatient Providers

ILI is defined as a fever of at least 100°F plus either a cough or a sore throat. In weeks when a relatively low number of enrolled facilities report data, the ILI proportion may not be as representative of Arizona activity as for other weeks. The state ILI baseline is 1.1% and the epidemic threshold is 1.9%*.

| | Week 38 | Week 39 |
|--|-----------------|-----------------|
| Proportion of patient visits to sentinel providers for ILI | 1.3% | 0.7% |
| Comparison to epidemic threshold* | Below threshold | Below threshold |
| Intensity level (see definitions at the end of report) | Minimal | Minimal |



*Note: The baseline is defined as the mean of the state ILI% in weeks in the 2011-2014 flu seasons in which two or more consecutive weeks each accounted for less than 2% of the season's total number of specimens testing positive for influenza at the Arizona State Public Health Laboratory. The epidemic threshold is defined as the mean plus two standard deviations.

Glossary of Key Terms:

2015-2016 Influenza Season – The season is defined by surveillance weeks. The first day of the 2015-2016 influenza season was October 4th, 2015, or week 40 and the 2015-2016 surveillance season will continue through October 1st, 2016, or week 39.

Regions – Regions in Arizona are defined by county: Central (Gila, Maricopa, Pinal); Northern (Apache, Coconino, Navajo, Yavapai); Southern (Cochise, Graham, Greenlee, Pima, Santa Cruz); Western (La Paz, Mohave, Yuma)

Activity Levels: Indicator of the geographic spread of influenza activity, reported to CDC by all states each week.

Widespread: Increased influenza-like illness from sentinel providers (ILI) in three or more regions and large numbers of laboratory-confirmed influenza cases in those regions.

Regional: Increased ILI in two regions and elevated numbers of laboratory-confirmed influenza cases in those regions.

Local: Increased ILI in one region and elevated numbers of laboratory-confirmed influenza cases in that region.

Sporadic: No increase in ILI activity and only isolated laboratory-confirmed influenza cases.

No Activity: No increase in ILI activity and no laboratory-confirmed influenza cases.

Intensity Levels: Intensity levels are based on the percent of outpatient visits in a state due to ILI and are compared to the average percent of ILI visits that occur during spring and fall weeks with little or no influenza virus circulation. Intensity levels range from minimal, corresponding to ILI activity from outpatient clinics being below the average, to intense, which would correspond to ILI activity from outpatient clinics being much higher than average.