

Position child and instruct parent in proper restraint. **DO NOT** attempt to immunize an uncontrolled child.
 If injecting multiple vaccines in same site, separate by 1 to 2 inches. Use no more than 2 vaccines in one muscle.

VACCINE	ROUTE	SITE
ROTAVIRUS	Oral	
e-IPV (enhanced IPV)	Subq.	Left arm Inject between shoulder and elbow in the back of the arm
DTaP/DT Pediarix & Pentacel Td – adult use	IM	Left Not Walking: Vastus lateralis, Antero lateral Thigh Walking: Deltoid
VARICELLA	Subq.	Left arm Inject between shoulder and elbow in the back of the arm
HEPATITIS A	IM	Right Deltoid If receiving multiple vaccines, use Left Deltoid
Hib Comvax- Hib/HepB	IM	Right Not Walking: Vastus lateralis, Antero lateral Thigh <u>Not Walking:</u> If receiving influenza, put Hib on Left Walking: Deltoid
HEPATITIS B	IM	Right Not Walking: Vastus lateralis, Antero lateral Thigh Walking: Deltoid
PCV 7 Pneumococcal Conjugate	IM	Right Not Walking: <u>Vastus lateralis, Antero lateral Thigh</u> If receiving multiple vaccines, no DTaP/DT - use Left thigh. Walking: <u>Deltoid</u> If receiving multiple vaccines, no DTaP/DT - use Left Deltoid for PCV7
Influenza	IM	Right Not Walking: <u>Vastus lateralis, Antero lateral Thigh</u> Walking: <u>Deltoid</u>
MMR / MMRV	Subq.	Right arm Inject between shoulder and elbow in the back of the arm

Subcutaneous injection use 5/8-inch, 25 gauge needle. Intramuscular injection use 1" to 1-1/2", 22 to 25 gauge needle.

Dispose of syringes and empty vaccine vials into red sharp containers.

MARICOPA COUNTY DEPARTMENT OF PUBLIC HEALTH - COMMUNITY HEALTH NURSING October 2008

