

**Children 11 Years and Older Need These Immunizations to Attend School in Arizona
2009-2010 School Year**

Immunizations	1 st Dose	2 nd Dose	3 rd Dose	4 th Dose	5 th Dose
<u>(DTaP/DTP) Diphtheria, Tetanus & Pertussis, (DT) Diphtheria & Tetanus, (Td) Tetanus & Diphtheria</u>	Required	Required	Required	Required if dose #1 was given before 12 months of age.	
An interval of at least 4 weeks is required between the first doses of DTaP/DTP/DT/Td . An interval of 6 months is required between the last two doses.					
<u>(Tdap) Tetanus, Diphtheria & Pertussis</u>	Required for 6 th and 7 th grade students who are 11 years and older and have had 5 years pass since their last DTaP/DTP//Td dose.				
Students 11 years and older in all grades are required to receive one dose of Tdap if 10 years have passed since their last dose of DTaP/DTP/DT/Td, or if they are in the process of receiving their first 3 doses of tetanus/diphtheria vaccine. Students who have received 1 dose of Tdap will need a Td booster after 10 years.					
<u>(MV) Meningococcal Vaccine</u>	Required for 6 th and 7 th grade students who are 11 years and older.				
A dose of MV is recommended at 11 years of age. Because MV is licensed for use in children 2 years and older, doses given between 2 and 11 years of age are valid and acceptable.					
<u>(IPV/OPV) Polio</u>	Required	Required	Required	Not required if dose #3 was given at 4+years. 4 doses meet the requirement even if all were given in the first year of life.	
An interval of at least 4 weeks is required between all Polio doses.					
<u>(MMR) Measles, Mumps, & Rubella</u>	Required	Required			
MMR #1 is due at 12-15 months of age. MMR #2 may be given 1 month after MMR #1 . MMR #2 is required for grades K-12.					
<u>(Hep B) Hepatitis B</u>	Required	Required	Required. Minimum age for final Hep B dose is 24 weeks.	Required only if dose #3 was given before 24 weeks of age.	
Hep B #1 may be given at birth. Dose #2 is due 4-8 weeks after dose #1. Dose #3 is due 2-5 months after dose #2, at least 4 months after dose #1 and no sooner than 24 weeks of age. A 2-dose adolescent series may be given to children 11-15 years of age over a 4-6 month period.					
<u>Varicella (Chicken Pox)</u>	Required	Required if dose #1 was given at 13+ years of age.			

In the 2009-2010 school year, children enrolled in preschool, kindergarten and grades 1, 2, 3, 4, 5, 7, 8, 9, 10 and 11 are required to be immunized against varicella (chicken pox) or have a history of chicken pox disease. Parental recall of chicken pox is acceptable. One dose of varicella vaccine meets the requirement for all children except those who receive varicella #1 at 13+ years of age. Children who receive the 1st dose of varicella vaccine at 13+ years of age need 2 doses, spaced at least 1 month apart.
Note: If MMR and Varicella vaccine doses were not given on the same day, they must be given at least 28 days apart.

