


# 2024 - 2025 PEDIATRIC INFLUENZA VACCINES

2024-2025 flu vaccines are all trivalent presentations

**\* Indicates this presentation is available to order through the Vaccines for Children (VFC) program.**


## 6 Months & Older




**Flucelvax® \***  
**Seqirus USA, Inc**  
0.5 mL single-dose syringe, 10 pack  
NDC 70461-0654-03



**Flucelvax® \***  
**Seqirus USA, Inc**  
5mL multi-dose vial, One 10-dose pack  
NDC 70461-0554-10




**Fluzone® \***  
**Sanofi Pasteur**  
0.5 mL single-dose syringe, 10 pack  
NDC 49281-0424-50



**Afluria®**  
**Seqirus USA, Inc**  
5mL multi-dose vial, One 10-dose pack  
NDC 33332-0124-10



**Fluzone® \***  
**Sanofi Pasteur**  
5mL multi-dose vial, One 10-dose vial  
NDC 49281-0641-15



**FluLaval® \***  
**GlaxoSmithKline Biologics**  
0.5 mL single-dose syringe, 10 pack  
NDC 19515-0810-52




# 2024 - 2025 PEDIATRIC INFLUENZA VACCINES

2024-2025 flu vaccines are all trivalent presentations


**\*** Indicates this presentation is available to order through the Vaccines for Children (VFC) program.

## 36 Months & Older



**Afluria® \***  
**Seqirus USA, Inc**  
0.5mL single dose syringe, 10 pack  
NDC 33332-0024-03

## 2 to 49 Years



**Flumist® \***  
**AstraZeneca**  
0.2 mL single-dose sprayer, 10 pack  
NDC 66019-0311-10


**Bureau of Immunization Services**  
**150 N 18th Ave., Suite 260**  
**Phoenix, AZ 85007**  
**Phone: 602-364-3630**  
**Fax: 602-364-3276**



**ARIZONA DEPARTMENT  
OF HEALTH SERVICES**  
PREPAREDNESS

# 2024 - 2025 PEDIATRIC INFLUENZA VACCINES

## Guide for Determining the Number of Doses of Influenza Vaccine to Give to Children Age 6 Months Through 8 Years



\*If a child previously received two doses of flu vaccine (at any time), they only need one dose of flu vaccine this season.

**Note:** If an 8-year-old patient needs two (2) doses of influenza vaccine this season, they should receive two (2) doses even if they turn 9 years old before the second dose is administered.

### Coadministration of COVID-19 and Influenza Vaccines

Providers can administer COVID-19 and influenza vaccines on the same day or coadminister at any time interval. There may be logistical advantages to administering COVID-19 and influenza vaccines on the same day, which providers may encourage to patients. There are no safety concerns for coadministration.

#### When deciding whether to coadminister another vaccine(s) with COVID-19 vaccine, consider:

- If the patient is behind OR if the patient is at risk of becoming behind on recommended vaccines
- The patient's risk of vaccine-preventable disease
- The reactogenicity profile of the vaccines
- The likelihood of avoiding a missed opportunity to vaccinate

#### Best practices for multiple injections include:

- Label each syringe with the name and the dosage (amount) of the vaccine, lot number, the initials of the preparer, and the exact beyond-use time, if applicable
- Separate injection sites by 1 inch or more, if possible
- Administer the COVID-19 vaccine and vaccines that may be more likely to cause a local reaction (i.e., tetanus-containing vaccines) in different limbs, if possible