




2026-2027 Pediatric Influenza Vaccines


6 Months & Older

Vaccine Packaging	Vaccine Name	Manufacturer	Presentation	Package Size	NDC	Available to order through VFC program?
	Flucelvax®	Seqirus USA, Inc	0.5 mL single-dose syringe	10 Pack	70461-0656-03	YES
	Fluzone®	Sanofi Pasteur	0.5 mL single-dose syringe	10 Pack	49281-0426-50	YES
	Fluarix®	GlaxoSmithKline Biologics	0.5mL single dose syringe	10 pack	58160-0725-52	YES
	FluLaval®	GlaxoSmithKline Biologics	0.5 mL single-dose syringe	10 Pack	19515-0822-52	YES

9 Years & Older

Vaccine Packaging	Vaccine Name	Manufacturer	Presentation	Package Size	NDC	Available to order through VFC program?
	Flublok®	Sanofi Pasteur	0.5 mL single-dose syringe	10 Pack	49281-0726-10	YES

2 to 49 Years

Vaccine Packaging	Vaccine Name	Manufacturer	Presentation	Package Size	NDC	Available to order through VFC program?
	Flumist®	AstraZeneca	0.2 mL single-dose sprayer	10 Pack	66019-0113-10	YES




For vaccines given
within City of Phoenix,
call 602-364-3899 or email
MCPIP@maricopa.gov



For vaccines given
outside City of Phoenix, call
602-364-3642 or email
ArizonaVFC@azdhs.gov

2026-2027 Pediatric Influenza Vaccines

Guide for Determining the Number of Doses of Influenza Vaccine to Give to Children Age 6 Months Through 8 Years



*If a child previously received two doses of flu vaccine (at any time), they only need one dose of flu vaccine this season.

Note: If an 8-year-old patient needs two (2) doses of influenza vaccine this season, they should receive two (2) doses even if they turn 9 years old before the second dose is administered.

Coadministration of COVID-19 and Influenza Vaccines

Providers can administer COVID-19 and influenza vaccines on the same day or coadminister at any time interval. There may be logistical advantages to administering COVID-19 and influenza vaccines on the same day, which providers may encourage to patients. There are no safety concerns for coadministration.

When deciding whether to coadminister another vaccine(s) with COVID-19 vaccine, consider:

- If the patient is behind OR if the patient is at risk of becoming behind on recommended vaccines
- The patient's risk of vaccine-preventable disease
- The reactogenicity profile of the vaccines
- The likelihood of avoiding a missed opportunity to vaccinate

Best practices for multiple injections include:

- Label each syringe with the name and the dosage (amount) of the vaccine, lot number, the initials of the preparer, and the exact beyond-use time, if applicable
- Separate injection sites by 1 inch or more, if possible
- Administer the COVID-19 vaccine and vaccines that may be more likely to cause a local reaction (for example, tetanus-containing vaccines) in different limbs, if possible