Healthcare Provider Office Information

We believe that getting vaccinated is a critical step in protecting your health. Vaccines can help prevent common diseases that can be serious and costly for you or your loved ones.

Each year, thousands of adults in America suffer serious health problems (and some even die) from diseases they could be vaccinated against like whooping cough, hepatitis A and B, flu and pneumococcal diseases, and shingles. Older adults and those with chronic health conditions are at increased risk for complication from certain diseases.

Together, let's take an active role in helping you and your loved ones stay healthy. To learn more about vaccines for adults, visit www.cdc.gov/vaccines/adults or take a quick vaccine quiz at www.cdc.gov/vaccines/adultquiz.

Please take a moment to fill out the questionnaire below to help us determine which vaccines may be recommended for you based on your specific health status, age, and lifestyle. Keep in mind that this list may not include every vaccine you need.

Check all that apply to you	Let's discuss these recommended vaccines
□ I am 19 years or older	Seasonal Flu (Influenza) vaccine every year Tetanus (Td) vaccine every 10 years One time dose of whooping cough (Tdap) vaccine for all adults who have never received Tdap vaccine PREGNANT WOMEN SHOULD GET A TDAP VACCINE DURING EACH PREGNANCY
☐ I am 60 years or older	Shingles (Zoster) vaccine*
☐ I am 65 years or older	Both types of pneumococcal vaccines (one dose of conjugate first, then one dose of polysaccharide 6-12 months later)
☐ I didn't receive the Human papillomavirus (HPV) vaccine series as a child	 HPV vaccine series (3 dose series) Female age 26 or younger Male age 21 or younger Male age 22-26 who has sex with men, who has a weakened immune system, or who has HIV
□ I was born in the US in 1957 or after and don't have immunity against measles, mumps, and rubella	Measles, mumps, rubella (MMR) vaccine* (one dose)
☐ I was born in the US in 1980 or after and don't have immunity against chickenpox	Varicella "chickenpox" vaccine*
☐ I am a healthcare worker	 Hepatitis B vaccine series Measles, mumps, rubella (MMR) vaccine* Varicella "chickenpox" vaccine*
☐ I have heart disease, asthma or chronic lung disease	Pneumococcal polysaccharide vaccine



Check all that apply to you	Let's discuss these recommended vaccines
☐ I have type 1 or type 2 diabetes	Hepatitis B vaccine series Pneumococcal polysaccharide vaccine
☐ I have a weakened immune system	 Both types of pneumococcal vaccines (one dose of conjugate first, then one dose of polysaccharide ≥8 weeks later) HPV vaccine series (if 26 years of age or younger and not previously vaccinated) Hib vaccine (post-hematopoietic stem cell transplant only)
□ I have HIV	 Hepatitis B vaccine series Both types of pneumococcal vaccines (one dose of conjugate first, then one dose of polysaccharide ≥8 weeks later) HPV vaccine series (if 26 years of age or younger and not previously vaccinated)
☐ I have chronic liver disease	 Hepatitis A vaccine series Hepatitis B vaccine series Pneumococcal polysaccharide vaccine
☐ I do not have a spleen or my spleen does not work well	 Hib vaccine Meningococcal vaccine Both types of pneumococcal vaccines (one dose of conjugate first, then one dose of polysaccharide ≥8 weeks later)
☐ I am a man who has sex with men	 Hepatitis A vaccine series Hepatitis B vaccine series HPV vaccine series (if 26 years of age or younger and not previously vaccinated)
☐ I am a laboratory worker and may be routinely exposed to isolates of Neisseria meningitidis, or specimens potentially containing hepatitis A or hepatitis B virus	 Hepatitis A vaccine series Hepatitis B vaccine series Meningococcal vaccine
☐ I am a college freshman living in a residence hall	Meningococcal vaccine Measles, mumps, rubella (MMR) vaccine*
☐ I am planning to travel out of the U.S.	Talk to your healthcare professional to learn which vaccines you may need based on locations of travel.

^{*}This is a live vaccine and should not be given to people who have a very weakened immune system, including those with a CD4 count less than 200, or to pregnant women.

