Normal Growth Patterns

Babies grow a lot in the first year of life. Your baby will triple their body weight by the time they are 12 months old. Breastfeeding moms may wonder if their baby is getting enough to eat because they cannot actually see milk going into their baby's mouth. One of the best ways to know that <u>your baby is getting enough</u> is by the number of wet and poopy diapers and their weight gain.

Wet Diapers

You can expect to see one wet diaper and one poopy diaper in the first 24 hours of life.

Then, you can expect another wet diaper for every 24 hours of life until day 5 to 7.

By day 5 to 7, your baby should have about 6 to 8 very wet diapers every 24 hours.

Poopy Diapers

Your baby will have a meconium diaper (black sticky poop) for the first two days and it will change to yellow seedy poop by day 5.

It is normal for a baby to poop every time they breastfeed, but not all babies do this.

The number of yellow poopy diapers may be different for each baby because it depends on how they are eating.



- During a growth spurt, your baby grows faster than usual and may suddenly eat more often. It is common for a mother to question her milk supply during a growth spurt.
- During this time, your baby will often show signs of hunger, like every hour or more. You may feel like you are not making enough milk or that your baby is not getting all he needs. You may think that you need to give formula or solid foods (supplement) or that it is time to stop breastfeeding. Be patient with yourself and your baby.
- Growth spurts only last 2 to 3 days. It is nature's way of increasing your milk supply for the future. Contact a lactation consultant if the growth spurt is lasting longer, as this may be a sign that your baby is having trouble getting milk from your breast.

Growth Spurts

A lot of growth happens in short quick bursts that can happen any time, but usually happen at...

- 3 weeks
- 6 weeks
- 8 weeks
- 3 months
- 6 months
- Your milk production depends on several things, including how often your baby breastfeeds. The more often your baby breastfeeds well, the more milk you will make.
- Giving formula or solid foods to your baby during a growth spurt may affect your milk supply in the future. Before supplementing, speak to a lactation consultant, your WIC counselor, or call the 24-Hour Breastfeeding Hotline at 1-800-833-4642.

Also available in Spanish.





24-Hour Breastfeeding Hotline 1-800-833-4642

This institution is an equal opportunity provider.
Arizona Department of Health Services 2024

Infant Growth and Behavior Patterns Quick Guide

| Months | Months | Months | Months |
|--------|--------|--------|--------|
| | | | |

| | Happy Day Days Days Weeks Months Months Months Months | | | | | | | Months | | |
|----------|---|--|---|---|---|---|---|--|-------------|---|
| | Happy Birthday | Day 2 | Days 3-4 | Days 5-11 | 2-4 | 1-4 | 5-6 | 6-8 | 9-12 | 13-15 |
| Feedings | Alert after first feed, may be sleepy the rest of the day. Provide lots of skin to skin care. | More alert, feeds every 2-4 hours. Lots of extra feedings during the second night. | 8-12 feeds in a 24-hour period (the feedings may not be evenly spaced). Look for hunger cues. | 8-12 feeds in a 24-hour period (the feedings may not be evenly spaced). Look for hunger cues. | 8-12 feeds in a 24-hour period (the feedings may not be evenly spaced). Look for hunger cues. | 8-12 feeds in a 24-hour period (the feedings may not be evenly spaced). Look for hunger cues. | 8-12 feeds in a 24-hour period (the feedings may not be evenly spaced). Look for hunger cues. | Time and nu breastfeedir after introdu foods. | ng may vary | May be once at night or several times throughout the day. |
| Diapers | Minimum 1 wet and 1 sticky black poop (meconium) | Minimum 2 poop and 2 wet | Minimum 3-4 wet, 3-4 poop. Poop starting to turn green and pasty. | Minimum 6+ wet, 3+ poop. Poop should be yellow, seedy and runny. | Minimum 6+ wet, 3+ poop. Poop should be yellow, seedy and runny. | After 6 weeks of age, pooping patterns can change and vary between babies. | | Introduction to solids can change the number, smell, and texture of poops. Your baby may not have a poop everyday. | | |
| Weight | Birth weight | Loses about 7% of their birth weight, which is normal. | Weight loss is small as baby begins to regain weight. | Weight gain of 0.5-1 ounces a day is normal. | Baby should be back up to birth weight at 2 weeks. Normal weight gain is 4-7 ounces a week. | Baby should be gaining 4-7 ounces per week after reaching his birth weight. | Baby should be gaining 4-5 ounces per week. | Baby should 2-4 ounces p | | Most toddlers gain 2 ounces per week. |

The following information is a guideline that can be followed, but the information should not replace your health care provider's recommendations. This can be used to have conversations with your baby's doctor. If you have specific concerns about your baby's growth or breastfeeding, please speak with your health care provider. Also, you can call the 24-Hour Breastfeeding Hotline at 1-800-833-4642 for any breastfeeding or baby behavior questions.